

# Corn Dust 2015

Chia Lin  
Reed Johnson



THE OHIO STATE UNIVERSITY

COLLEGE OF FOOD, AGRICULTURAL,  
AND ENVIRONMENTAL SCIENCES




A



B






SE

22-Apr-15

WD27.9mm 15.0kV x1.0k 50um




Tier I  
Level of Concern  
(LOC) for foliar  
application of  
clothianidin =  
 $0.7 \text{ ng /cm}^2$

# Field Experiments




# Study sites



FSR



Radius = 2 km





## Apiary setup:

Tracked 2 - 6 actively foraging,  
queen-right colonies per site

Monitored hive conditions: queen  
status, colony growth, honey,  
pollen, and mite loads

Pollen traps were installed on 2 -  
4 additional hives for pollen  
sampling



Pollen trap "ON"




Pollen trap "OFF"








Use “Drop Zone Dead Bee Traps” to collect dead bees







# Hive monitoring and maintenance

## Detailed inspections

- pre-planting (April 27-30)
- post-planting I (May 20-22)
- post-planting II (June 19 - 24)
- post-planting III (August 14 - 22)

## Honey harvest:

- June/September

## Mite treatment:

- Apivar: before experiment
- Formic acid: June, September
- Oxalic acid: November - December

## Feeding (as needed):

- November - February





# Quantifying hive parameters

## “Box crack” inspections:

- seams of bees
- mite count (sugar shake)

## Frame area inspections:

- bees
- capped brood
- open brood
- honey
- pollen
- empty drawn comb
- undrawn foundation
- drone brood
- queen cells




# Quantifying hive parameters

“Box crack” inspections:



- seams of bees
- mite count (sugar shake)

Frame area inspections:

- bees
- capped brood
- open brood
- honey
- pollen
- empty drawn comb
- undrawn foundation
- drone brood
- queen cells




# Trends of increasing neonic concentration in pollen with more corn in the surrounding landscape




# Hive-level effects: worker bees

## Seams of bees




## Area of bees




# Hive-level effects: brood

## Capped brood




## Open brood




# Hive-level effects: food storage


## Pollen



## Nectar








Khoury et al., 2013

# Closed Nuc Experiments



Bee Brief 4-frame nuc box

**Assembled nucs > 4 weeks before experiment:**

- 1 frame of honey
- 1 frame of capped brood
- 1 - 2 frames of bees
- ripe queen cells supplied by a local breeder



## Three days before experiment:

- Identified 6 healthy nucs with actively laying new queens
- Caged the queen to ensure no egg was present when the trial began





## Treatments:



**150 g pollen + 20ml 50% sugar syrup**

**Control:** pollen with ~ 0 ppb residue


**Low exposure:** pollen with 10 - 20 ppb clo. + thi.


**High exposure:** pollen with > 20 ppb clo. + thi. (max = 107 ppb)

**Positive control:** pollen with ~ 0 ppb residue, syrup spiked with clothianidin (final concentration = 400 ppb)



Biotrix microplates





Foundation  
+ pollen

Brood

Empty  
comb

Foundation



Closed nucs were kept indoor during trials (5 trials total)  
Opened on days 4, 8, and 12 for sampling and recording data





## Sampling:

- pollen
- nurse bees
- mature larvae if available






# Mobile bee studio




## Photographing frame status

- each side of a frame is divided to 4 quarters, each assigned with a unique identifier barcode
- images of the quarters are re-assembled and frame contents are quantified using computer programs





150500510224  
Exp: 5 Nov 5 0.3  
Ex: 2 Nov 0 TOP






250400130224  
Exp: 4 Nov 1 013  
Fr: 12 Pa: 00 TOP






# Computer detection of cell location



# Automated Cell Scoring Software





Determine fate of each egg laid

